

MAXIMAL MULTIHOMOGENEITY OF ALGEBRAIC HYPERSURFACE SINGULARITIES

MATHIAS SCHULZE

ABSTRACT. From the degree zero part of logarithmic vector fields along an algebraic hypersurface singularity we identify the maximal multihomogeneity of a defining equation in form of a maximal algebraic torus in the embedded automorphism group. We show that all such maximal tori are conjugate and in one-to-one correspondence to maximal tori in the degree zero jet of the embedded automorphism group.

The result is motivated by Kyoji Saito's characterization of quasihomogeneity for isolated hypersurface singularities [Sai71] and extends its formal version [GS06, Thm.5.4] and a result of Hauser and Müller [HM89, Thm.4].

Let $\mathcal{P}$ be either $\mathcal{O} = \mathbb{C}\{x\}$, the ring of convergent power series in $x = x_1, \dots, x_n$, or $\hat{\mathcal{O}} = \mathbb{C}[[x]]$, the ring of formal power series in x and denote by $\mathfrak{m} = \langle x \rangle$ the maximal ideal in $\mathcal{O}$ or $\hat{\mathcal{O}}$. Let $\text{Aut}(\mathcal{P})$ be the automorphism group of $\mathcal{P}$ and let $\Delta(\mathcal{P}) = \mathfrak{m} \cdot \text{Der}(\mathcal{P}) = \{\delta \in \text{Der}(\mathcal{P}) \mid \delta(\mathfrak{m}) \subseteq \mathfrak{m}\}$ be the $\mathcal{P}$ -module of vector fields vanishing at the origin.

Let $0 \neq f \in \mathfrak{m}$ and $\text{Aut}_f = \{\varphi \in \text{Aut}(\mathcal{P}) \mid \varphi(f) \in \langle f \rangle\}$ the group of automorphisms preserving the ideal $\langle f \rangle$. In the convergent case, this is the group of automorphisms of $(\mathbb{C}^n, 0)$ preserving the hypersurface $\text{Var}(f)$. Our object of interest is the $\mathcal{P}$ -module of logarithmic vector fields $\text{Der}_f = \{\delta \in \text{Der}(\mathcal{P}) \mid \delta(f) \in \langle f \rangle\}$ introduced in [Sai80]. In the convergent case, this is the module of vector fields tangent to the smooth part of $\text{Var}(f)$. The module Der_f is unchanged if we assume f to be reduced. We shall further assume that $\text{Der}_f \subseteq \Delta(\mathcal{P})$. By Rossi's Theorem [Ros63, Cor.3.4], this means in the convergent case that the variety $\text{Var}(f)$ defined by f is not a product with a smooth factor. As remarked in [HM93, 2.Rem.(c)], Der_f is the Lie algebra of the infinite Lie group Aut_f in the convergent case.

For a fixed coordinate system, any derivation $\delta \in \text{Der}(\mathcal{P})$ can be decomposed into homogeneous components, $\delta = \sum_{i=-1}^{\infty} \delta_i$. Moreover, $\delta_0 = \sum_{i,j} a_{i,j} x_i \partial_{x_j}$ for some matrix $A = (a_{i,j})$ and we call δ_0 diagonal if A is diagonal. A derivation $\delta \in \Delta(\mathcal{P})$ is called nilpotent if A is nilpotent and semisimple if $\mathfrak{m}$ has a basis of eigenvectors of δ . For a fixed coordinate system, any $\delta \in \Delta(\mathcal{P})$ is a sum $\delta = \delta_S + \delta_N$ where $\delta_S = \delta_{S,0}$ and $\delta_N, 0$ are defined by the semisimple and nilpotent parts of the matrix A corresponding to δ_0 . In particular, $[\delta_S, \delta_{N,0}] = 0$.

In [GS06, Thm.5.4], K. Saito's construction [Sai71, §3] of the Poincaré–Dulac decomposition [AA88, Ch.3.§3.2] is generalized to Part a of the following theorem.

Date: November 8, 2006.

1991 *Mathematics Subject Classification.* 32S25, 17B15.

Key words and phrases. hypersurface singularity, multihomogeneity, logarithmic vector field.

The author is grateful to Michel Granger and Gerhard Pfister for helpful discussions and acknowledges partial support by the Humboldt foundation.

The statement a.5 is implicitly present in the arguments in [GS06]. We shall follow the outline of [HM89, Thm.4] to deduce Part b of Theorem 1 from Part a.

Theorem 1 (existence of maximal multihomogeneity). *Let*

- a. $f \in \hat{\mathcal{O}}$ or
- b. $f \in \mathcal{O}$ algebraic over $\mathbb{C}[x]$

and let $\delta_1, \dots, \delta_t \in \text{Der}_f$ with diagonal degree 0 part. Then there is an algebraic torus $T^s \subseteq \text{Aut}_f$ in the sense of [Mül86, 1.Def.ii)] with Lie algebra $\mathfrak{t}^s$ and suitable coordinates such that $T^s \subseteq \text{GL}_n(\mathbb{C})$. In these coordinates there is a basis $\sigma_1, \dots, \sigma_s$ of $\mathfrak{t}^s$ which extends to a minimal system of generators $\sigma_1, \dots, \sigma_s, \nu_1, \dots, \nu_r$ of Der_f and a choice of f with irreducible factors $f_1, \dots, f_m$ such that

- 1. σ_i is diagonal with eigenvalues in $\mathbb{Z}$,
- 2. ν_i is nilpotent,
- 3. $[\sigma_i, \nu_j] \in \mathbb{Z} \cdot \nu_j$,
- 4. $\sigma_i(f_j) \in \mathbb{Z} \cdot f_j$,
- 5. $(\delta_i)_0 \in \langle \sigma_1, \dots, \sigma_s \rangle_{\mathbb{C}}$, and
- 6. if $\delta \in \text{Der}_f$ with $[\sigma_i, \delta_0] = 0$ for all i then $\delta_S \in \langle \sigma_1, \dots, \sigma_s \rangle_{\mathbb{C}}$.

Proof. The statement a.5 holds by the construction in the proof of [GS06, Thm.5.4] using that the coordinate change in [GS06, Thm.5.3] is tangent to the identity.

By a.1 and a.4, there is a formal coordinate change $\bar{y}(x)$ such that $\bar{g}(x) = f(\bar{y}(x))$ fulfils $\sigma_i(\bar{g}) = \lambda_i \cdot \bar{g}$ where $\lambda_i \in \mathbb{Z}$ and $\sigma_i = \sum_{j=1}^n \sigma_{i,j} x_j \partial_{x_j}$ where $\sigma_{i,j} \in \mathbb{Z}$. We identify σ_i with its coefficient vector $(\sigma_{i,1}, \dots, \sigma_{i,n}) \in \mathbb{Z}^n$. By [Bor91, II.7.3 (2)], the the saturation

$$L = ((\mathbb{Z}^n)^\vee / (\mathbb{Z}^n / \sum_{i=1}^s \mathbb{Z} \sigma_i)^\vee)^\vee \cong \mathbb{Z}^s \subseteq \mathbb{Z}^n$$

of the lattice generated by $\sigma_1, \dots, \sigma_s$ defines an algebraic torus

$$G_m^s \cong T^s := \text{Spec}(\mathbb{C}[L]) \subseteq \text{Spec}(\mathbb{C}[\mathbb{Z}^n]) \subseteq \text{GL}_n(\mathbb{C}) \subseteq \text{Aut}(\mathcal{P})$$

with Lie algebra $\mathfrak{t}^s = \langle \sigma_1, \dots, \sigma_s \rangle$. Since the x -monomials are common eigenvectors for $x_1 \partial_1, \dots, x_n \partial_n$ with integer eigenvalues, we may assume $L = \sum_{i=1}^s \mathbb{Z} \sigma_i$ preserving the condition $\lambda_i \in \mathbb{Z}$. Then the σ -multihomogeneity of $\bar{g}$ of multidegree λ stated above translates to $\bar{g}$ being equivariant for $T^s \subseteq \text{GL}_n(\mathbb{C})$ and the character $T^s \rightarrow \text{Spec}(\mathbb{C}[\mathbb{Z}]) = \text{GL}_1(\mathbb{C})$ defined by $L \ni \sigma_i \mapsto \lambda_i \in \mathbb{Z}$. Now $\langle f \rangle_{\hat{\mathcal{O}}}$ is equivalent to the T^s -stable ideal $\langle \bar{g} \rangle_{\hat{\mathcal{O}}}$ and [HM89, Thm.2'] implies that also $\langle f \rangle_{\mathcal{O}}$ is equivalent to a T^s -stable ideal. This means that there is an analytic coordinate change $y(x)$ and a unit $u \in \mathcal{O}^*$ such that $g(x) = u(x) \cdot f(y(x))$ generates a T^s -stable ideal in $\mathcal{O}$. By abuse of notation we denote this g by f again. Then [Mül86, Hilfssatz 2] shows that $\langle f \rangle_{\mathcal{O}}$ has a T^s -equivariant generator which we may assume to be f . This proves b.1 and $\sigma_i(f) \in \mathbb{Z} \cdot f$ which is weaker than b.4.

To avoid the non-trivial [Mül86, Hilfssatz 2] and conclude b.4, one can argue as follows: By [Hum75, Thm.13.2], the T^s - and $\mathfrak{t}^s$ -stable subspaces in $\mathcal{O}/\mathfrak{m}^k$ coincide. Since ideals in $\mathcal{O}$ are closed in the $\mathfrak{m}$ -adic topology, this shows that $\langle f \rangle_{\mathcal{O}}$ is σ -stable. Then a multigraded version¹ of [SW73, 2.4] shows that $\langle f \rangle_{\mathcal{O}}$ has a T^s -equivariant generator. With $\langle f \rangle_{\mathcal{O}}$ also its minimal associated primes are σ -stable by [SW73, 2.5]. Again by [SW73, 2.4], these primes have T^s -equivariant generators $f_1, \dots, f_m$ and b.4 follows.

¹The arguments in [SW73, 2.2-4] give a more general correspondence of $(k^s, +)$ -graduations and sets of s simultaneously diagonalizable k -derivations on analytic k -algebras.

With f also its partial derivatives $\partial_{x_i}(f)$ are T^s -equivariant. Thus Der_f is T^s -stable since it can be considered as the projection of the syzygy module of $\partial_{x_1}(f), \dots, \partial_{x_n}(f), f$ to the first n components. Its T^s -sub-module $\mathfrak{t}^s = \langle \sigma_1, \dots, \sigma_s \rangle$ is the adjoint representation of T^s and hence trivial. By a module version² of [Mül86, Hilfssatz 2], Der_f has a minimal system of generators $\sigma_1, \dots, \sigma_s, \nu_1, \dots, \nu_r$ which spans over $\mathbb{C}$ a rational T^s -module. By the proof of [Mil06, Thm.9.13] the T^s -module $\langle \sigma_1, \dots, \sigma_s, \nu_1, \dots, \nu_r \rangle_{\mathbb{C}}$ can be diagonalized without changing the σ_i and b.3 follows. Like above, one can use a multigraded module version of [SW73, 2.4] instead.

The maximality property b.6 follows from its formal version a.6. Combined with b.3 it guarantees the existence of ν_i satisfying b.2: If the σ -multidegree of a ν_i is non-zero then it is nilpotent by [GS06, Lem.2.6], otherwise one can subtract its semisimple part which is a linear combination of the σ_i by b.6. $\square$

Remark 1.

(1) By [KPR75, 2.11], the Implicit Function Theorem holds in $\mathbb{C}\langle x \rangle$, the ring of algebraic power series. Thus the proof of [HM89, Thm.2'] works inside of $\mathbb{C}\langle x \rangle$ as well. However it is not clear if this ring is graded in the sense of [SW73, §1-2]. A positive answer would imply that Theorem 1.b even holds inside of $\mathbb{C}\langle x \rangle$.

(2) According to Michel Granger, the uniqueness of s in Theorem 1 can be deduced from the conjugacy of all Cartan subalgebras [Ser01, III.4.Thm.2] by showing that $\mathfrak{t}^s$ and $\mathfrak{n}_0 = \langle (\nu_j)_0 \mid \sigma_i(\nu_j) = 0 \rangle_{\mathbb{C}}$ span a Cartan subalgebra $\mathfrak{h}$ of $\mathfrak{g} = \text{Der}_f / (\text{Der}_f \cap \mathfrak{m}^2 \cdot \text{Der}(\mathcal{P}))$ where $\mathfrak{n}_0$ is the intersection of $\mathfrak{h}$ with the set of nilpotent elements in $\mathfrak{g}$.

We shall use results of Müller [Mül86] to prove a stronger statement than uniqueness of s . Consider the group morphisms $\pi_k : \text{Aut}(\mathcal{P}) \rightarrow \text{Aut}_k(\mathcal{P}) = \text{Aut}(\mathcal{P}/\mathfrak{m}^{k+2})$ and the Lie algebra morphisms $\pi_k : \Delta(\mathcal{P}) \rightarrow \Delta_k(\mathcal{P}) = \Delta(\mathcal{P})/(\mathfrak{m}^{k+1} \cdot \Delta(\mathcal{P}))$. Note that $\text{Aut}_k(\mathcal{P})$ is an algebraic group with Lie algebra $\Delta(\mathcal{P})$. Like in [Mül86, §2], one can use Artin's Approximation Theorem [Art68, Thm.1.2] to prove the first part of the following

Lemma 1. $\pi_k(\text{Aut}_f)$ is an algebraic group with Lie algebra $\pi_k(\text{Der}_f)$.

Proof. By exactness of completion $\text{Der}_f = \widehat{\text{Der}_f}$ where $\hat{f}$ denotes f considered in $\hat{\mathcal{O}}$. Thus $\pi_k(\text{Der}_f) = \pi_k(\text{Der}_{\hat{f}})$ and we may assume that $\mathcal{P} = \hat{\mathcal{O}}$. Consider the Lie algebra morphisms $\pi_k^m : \Delta_m(\mathcal{P}) \rightarrow \Delta_k(\mathcal{P})$ and denote by f_k the image of f in $\mathcal{O}/\mathfrak{m}^{k+1}$. For fixed k , the $\pi_k^m(\text{Der}_{f_m})$ form a decreasing sequence of subvector spaces in $\Delta_k(\mathcal{P})$ which implies $\pi_k^m(\text{Der}_{f_m}) = D_k$ for large m . Then $\pi_m^{m+1} : D_{m+1} \rightarrow D_m$ is surjective and hence $D_k = \pi_k(\text{Der}_f)$. In the proof of the first statement, we find $\pi_k^m(\text{Aut}_{f_m}) = \pi_k(\text{Aut}_f)$ for large m and Der_{f_m} is the Lie algebra of Aut_{f_m} . $\square$

Theorem 2 (uniqueness of maximal multihomogeneity). *The algebraic torus T^s in Theorem 1 is maximal in Aut_f and also $\pi_0(T^s) \subseteq \pi_0(\text{Aut}_f)$ is a maximal algebraic torus. All maximal algebraic tori in Aut_f and $\pi_0(\text{Aut}_f)$ are conjugate.*

Proof. First let $T^t \subseteq \pi_0(\text{Aut}_f)$ be an algebraic torus with Lie algebra $\mathfrak{t}^t$. If $\pi_0(T^s) \subseteq T^t$ then the Lie algebra of T^t consists of σ -homogeneous semisimple elements of

²The statement in [Mül86, Hilfssatz 2] holds more generally for any analytic sub-module of a free analytic module in the sense of [GR71].

multidegree 0 and hence $\pi_0(\mathfrak{t}^s) = \mathfrak{t}^t$ by Theorem 1.6. By Lemma 1 and [Mil06, Cor.13.11], this implies $\pi_0(T^s) = T^t$ and hence $\pi_0(T^s)$ is maximal.

Now let $T^t \subseteq \text{Aut}_f$ be an algebraic torus in the sense of [Mül86, 1.Def.ii)] with Lie algebra $\mathfrak{t}^t$. By definition, $\pi_k(T^t) \subseteq \pi_k(\text{Aut}_f) \subseteq \text{Aut}_k(\mathcal{P})$ defines a rational representation of T^t which is diagonalizable by [Mil06, Thm.9.13]. Then Cartan's Uniqueness Theorem holds for T^t by [Kau67, Satz] and π_0 restricted to T^t is injective. This shows that maximality of $\pi_0(T^s)$ implies maximality of T^s .

By [Hum75, VIII.21.3.Cor.A], all maximal tori of $\pi_0(\text{Aut}_f)$ are conjugate. An embedded version³ of [Mül86, Satz 2] shows that a conjugacy in $\pi_0(\text{Aut}_f)$ of algebraic tori in Aut_f lifts to a conjugacy in Aut_f . $\square$

Corollary 1 (lifting of maximal tori). $\pi_0 : \text{Aut}_f \rightarrow \pi_0(\text{Aut}_f)$ defines a bijection of algebraic tori.

REFERENCES

- [AA88] D. V. Anosov and V. I. Arnol'd (eds.), *Dynamical systems. I*, Encyclopaedia of Mathematical Sciences, vol. 1, Springer-Verlag, Berlin, 1988, Ordinary differential equations and smooth dynamical systems, Translated from the Russian [MR0823488 (86i:58037)]. MR MR970793 (89g:58060)
- [Art68] M. Artin, *On the solutions of analytic equations*, Invent. Math. **5** (1968), 277–291. MR MR0232018 (38 #344)
- [Bor91] Armand Borel, *Linear algebraic groups*, second ed., Graduate Texts in Mathematics, vol. 126, Springer-Verlag, New York, 1991. MR MR1102012 (92d:20001)
- [GR71] H. Grauert and R. Remmert, *Analytische Stellenalgebren*, Springer-Verlag, Berlin, 1971, Unter Mitarbeit von O. Riemenschneider, Die Grundlehren der mathematischen Wissenschaften, Band 176. MR MR0316742 (47 #5290)
- [GS06] Michel Granger and Mathias Schulze, *On the formal structure of logarithmic vector fields*, Comp. Math. **142** (2006), 765–778.
- [HM89] Herwig Hauser and Gerd Müller, *Algebraic singularities have maximal reductive automorphism groups*, Nagoya Math. J. **113** (1989), 181–186. MR MR986444 (90c:32041)
- [HM93] ———, *Affine varieties and Lie algebras of vector fields*, Manuscripta Math. **80** (1993), no. 3, 309–337. MR MR1240653 (94j:17025)
- [Hum75] James E. Humphreys, *Linear algebraic groups*, Springer-Verlag, New York, 1975, Graduate Texts in Mathematics, No. 21. MR MR0396773 (53 #633)
- [Kau67] Wilhelm Kaup, *Einige Bemerkungen über Automorphismengruppen von Stellenringen*, Bayer. Akad. Wiss. Math.-Natur. Kl. S.-B. **1967** (1967), no. abt. II, 43–50 (1968). MR MR0235064 (38 #3376)
- [KPR75] H. Kurke, G. Pfister, and M. Roczen, *Henselsche Ringe und algebraische Geometrie*, VEB Deutscher Verlag der Wissenschaften, Berlin, 1975, Mathematische Monographien, Band II. MR MR0491694 (58 #10899)
- [Mil06] J.S. Milne, *Algebraic groups and arithmetic groups*, 2006, <http://www.jmilne.org/math/CourseNotes/aag.html>.
- [Mül86] Gerd Müller, *Reduktive Automorphismengruppen analytischer $\mathbf{C}$ -Algebren*, J. Reine Angew. Math. **364** (1986), 26–34. MR MR817635 (88d:32041)
- [Ros63] Hugo Rossi, *Vector fields on analytic spaces*, Ann. of Math. (2) **78** (1963), 455–467. MR MR0162973 (29 #277)
- [Sai71] Kyoji Saito, *Quasihomogene isolierte Singularitäten von Hyperflächen*, Invent. Math. **14** (1971), 123–142. MR MR0294699 (45 #3767)
- [Sai80] ———, *Theory of logarithmic differential forms and logarithmic vector fields*, J. Fac. Sci. Univ. Tokyo Sect. IA Math. **27** (1980), no. 2, 265–291. MR MR586450 (83h:32023)
- [Ser01] Jean-Pierre Serre, *Complex semisimple Lie algebras*, Springer Monographs in Mathematics, Springer-Verlag, Berlin, 2001, Translated from the French by G. A. Jones, Reprint of the 1987 edition. MR MR1808366 (2001h:17001)

³In the first paragraph of [Mül86, §5], the proof of [Mül86, Satz 2] is reduced to the case of embedded automorphism groups using [Mül86, Satz 6].

- [SW73] Günter Scheja and Hartmut Wiebe, *Über Derivationen von lokalen analytischen Algebren*, Symposia Mathematica, Vol. XI (Convegno di Algebra Commutativa, INDAM, Rome, 1971), Academic Press, London, 1973, pp. 161–192. MR MR0338461 (49 #3225)

MATHIAS SCHULZE, DEPARTMENT OF MATHEMATICS, OKLAHOMA STATE UNIVERSITY, 401 MSCS,
STILLWATER, OK 74078, USA

E-mail address: `mschulze@math.okstate.edu`